

Tonisity Px The Quickstart Guide

Mixing Instructions:

- Mix 310g (2 level 155g scoops) of Tonisity Px with 10 L of fresh water to make a 3% liquid Tonisity Px solution.
- For best results, add Tonisity Px into water and stir until the powder is dissolved.
- Make fresh feed daily and use clean feeder pans for delivery.



For accurate mixing, use the Tonisity Px scoop to measure the right volume of powder.



Important Points

- Always use fresh potable water for mixing the Tonisity Px solution.
- Mix only enough solution for one feeding at a time.
- Add powder to water (about 25°C) and stir gently until fully dissolved by using a wire whisk or similar tool.
- Do not use hot water (above 40°C).
- Wash, rinse and sanitize utensils after every feed.



TIP:

To encourage consumption of the solution at Day 2, dip the nose of a few piglets into the solution. They will teach the other ones how to drink.

Suckling:

From Day 2-8 of life, give 500ml of Tonisity Px 3% solution per litter per day (or 40ml/piglet day).

Creep Feed Introduction:

For first 3 days of creep feeding, moisten creep feed with approx. 20ml solution for every 100g of feed



Weaning Transition:

For 3 days pre-weaning give 500ml of solution/litter/day or 40ml/piglet/day on top of creep feed. For 3 days post-weaning give up to 100ml solution/piglet/day on top of creep feed.

Transport Hydration:

For newly-weaned pigs, follow the weaning transition protocol. For grow-finish pigs give 100ml of solution per 10kg liveweight for 1-2 days before and after transport.

Precautionary Measures

- Make sure piglets have access to clean, fresh water at all times.
- Before feeding, ensure that pans are clean.
- Position the feeding pan in an area where piglets can easily access (near the nest or drinkers) and not in a cold or dirty corner.
- If product is left in pans after half a day, give to the sows.
- Chlorine treatment of groundwater can be irregular. A high content of chlorine in water will negatively affect the taste of Tonisity Px solution.

Storage and Handling

- Storage: Store the Tonisity Px powder below 30°C (room temperature) in clean, dry conditions.
- Do not expose to direct sunlight or moisture.
- To maintain freshness, keep partially used bags closed tightly after opening.

Cautionary Practice

Directions for use must be carefully followed.

Tonisity Px is not for human consumption.

Consult the Tonisity website www.tonisity.com or contact our local representative for further information.

16 Fitzwilliam Place, Dublin 2, Ireland **T:** +353 (0)1 902 0026 **E:** orders@tonisity.com **W:** tonisity.com

Tonisity Px[™] is a trademark of Tonisity Int. Ltd. ©2018 Tonisity International Ltd. PX-18-03-METRIC

Tonisity PX is an isotonic nutritional supplement designed specifically for pigs. It is not a drug and it does not contain ingredients with drug-like properties. It is not intended to diagnose, treat, cure, or prevent any disease. Any observed differences in performance are due to the nutritional and hydration properties of Tonisity PX.